

tPA/PAI-1 depleted plasma

REF 273

Description

Lyophilized normal citrated human plasma depleted of tPA and PAI-1 via immunoaffinity chromatography, using columns of antibodies specific for human tPA and human PAI-1 immobilized to agarose beads.

For Research Use Only.

Presentation

Screw-capped glass vial containing the equivalent of 1.0 mL plasma. The plasma contains 20 mM HEPES buffer.

Reconstitution

Add 1.0 mL of filtered deionized/distilled water.

Specification

<u>Test</u>	<u>Assayed Values</u>
PT	XX.X sec
APTT	XX.X sec
Fibrinogen	X.XX g/L
tPA	< 0.5 ng/mL
PAI-1	< 0.5 ng/mL

Reagent Storage and Handling

Store at 2° - 8°C. Reconstituted plasma should be held on melting ice for the duration of the testing. The material should be used within 4 hours of reconstitution.

WARNING

The source material for this reagent has been found to be non-reactive for Hepatitis B Surface Antigen (HBsAg), Hepatitis C Virus (HCV) and Human Immunodeficiency Virus Type 1 and Type 2 (HIV-1, HIV-2) using FDA cleared methods. As no known test method can provide complete assurance that products derived from human blood will not transmit HBsAg, HCV, HIV-1, HIV-2 or other blood-borne pathogens, this reagent should be handled as recommended for any potentially infectious human specimen.