

TFPI depleted plasma

REF 848DP

Description

Lyophilized normal citrated human plasma immunodepleted of tissue factor pathway inhibitor (TFPI) via immunoaffinity chromatography, using a column of antibody specific for human TFPI immobilized to agarose beads.

Presentation

Screw-capped glass vial containing the equivalent of 0.5 mL plasma. The plasma contains 20 mM HEPES buffer.

Reconstitution

Add 0.5 mL of filtered deionized/distilled water.

Reagent Storage and Handling

Store at 2° - 8°C. Reconstituted plasma should be held on melting ice for the duration of the testing. The material should be used within 4 hours of reconstitution.

Warnings and Precautions

For Research Use Only.

The source material for this reagent has been found to be non-reactive for Hepatitis B Surface Antigen (HBsAg), Hepatitis C Virus (HCV) and Human Immunodeficiency Virus Type 1 and Type 2 (HIV-1, HIV-2) using FDA cleared methods. As no known test method can provide complete assurance that products derived from human blood will not transmit HBsAg, HCV, HIV-1, HIV-2 or other blood-borne pathogens, this reagent should be handled as recommended for any potentially infectious human specimen.

Definition of Symbols

	Consult instructions for use
	Manufacturer
	Refer to Safety Data Sheet
	Temperature Limitation
LOT	Lot Number
	Expiration Date
REF	Catalog Number